


Theorem 1.1 [connection figure (1122 3344 5566 7788)]



(i) Let k, r, s, N be natural numbers, such that the 16 numbers

(#) $1, k+1, r+1, s+1, k+r+1, k+s+1, r+s+1, k+r+s+1,$
 $N-k-r-s, N-r-s, N-k-s, N-k-r, N-s, N-r, N-k, N$

are pairwise different and positive.

Then there exist 384 general 4×4 magic squares $M001, M002, \dots, M384$ with entries from (#), of connection figure (1122 3344 5566 7788) namely:

$$M001 = \begin{pmatrix} 1 & N & k+r+1 & N-k-r \\ k+s+1 & N-k-s & r+s+1 & N-r-s \\ N-k & k+1 & N-r & r+1 \\ N-s & s+1 & N-k-r-s & k+r+s+1 \end{pmatrix}, \quad M002 = \begin{pmatrix} 1 & N & N-k-s & k+s+1 \\ N-k-r & k+r+1 & r+s+1 & N-r-s \\ k+r+s+1 & N-k-r-s & N-r & r+1 \\ N-s & s+1 & k+1 & N-k \end{pmatrix},$$

$$M003 = \begin{pmatrix} 1 & N & N-r-s & r+s+1 \\ k+s+1 & N-k-s & N-k-r & k+r+1 \\ N-k & k+1 & k+r+s+1 & N-k-r-s \\ N-s & s+1 & r+1 & N-r \end{pmatrix}, \quad M004 = \begin{pmatrix} 1 & N & k+r+1 & N-k-r \\ N-r-s & r+s+1 & N-k-s & k+s+1 \\ N-k & k+1 & N-r & r+1 \\ k+r+s+1 & N-k-r-s & s+1 & N-s \end{pmatrix},$$

$M004$ till $M008$ are derived from $M001$ till $M004$ by simultaneous exchange of their rows 1 with 2, the rows 3 with 4, and then the columns 1 with 2 and the columns 3 with 4. $M009, M010, \dots, M016$ arise from $M001, \dots, M008$ by reflection at a horizontal axis, further the squares $M017$ till $M032$ are the mirror images of $M001, \dots, M016$ from reflection at a vertical axis. $M033, M034, \dots, M192$ come from $M001, \dots, M032$ by application of the 5 non-identical permutations of three objects to the triple (k, r, s) and finally, $M193, M194, \dots, M384$ are derived from $M001, \dots, M192$ by replacement of each entry x by $N+1-x$ (which has the same effect as simultaneous exchange of column 1 with column 2 and column 3 with column 4).

(ii) Let T be a symmetric subset of $\{1, \dots, N\}$ with 16 elements, containing 1 as an element and let M be a general 4×4 magic square of connection figure (1122 3344 5566 7788) with entries from T . Then there exists a triple (k, r, s) of natural numbers with $k < r < s$ such that T consists of the numbers (#). When 1 is a diagonal element of M , then M is one of the squares $M001, \dots, M192$; otherwise M is one of the squares $M193, \dots, M384$.

Proof

(i) can be verified easily. (ii) can be shown by solving the involved linear equations for M .

Definition 1

Let x_1, x_2, x_3, x_4 be natural numbers, such that $x_1 < x_2 < x_3 < x_4$ and $1+x_1+x_2+x_3+x_4=N$ and the 16 numbers

$$(*) \quad S(1)=1, S(2)=1+x_1, S(3)=1+x_2, S(4)=1+x_1+x_2, \\ S(5)=1+x_3, S(6)=1+x_1+x_3, S(7)=1+x_2+x_3, S(8)=1+x_1+x_2+x_3, \\ S(9)=1+x_4, S(10)=1+x_1+x_4, S(11)=1+x_2+x_4, S(12)=1+x_1+x_2+x_4, \\ S(13)=1+x_3+x_4, S(14)=1+x_1+x_3+x_4, S(15)=1+x_2+x_3+x_4, S(16)=N$$

are pairwise different.

Call a set of 4 numbers $\{S(i_1), S(i_2), S(i_3), S(i_4)\}$ with $S(i_1)+S(i_2)+S(i_3)+S(i_4)=2(N+1)$

"correct", if 1, x_1, x_2, x_3 , and x_4 each occur exactly 2 times, when the 4 numbers are represented by 1, x_1, x_2, x_3, x_4 .

Theorem 1.2

Let x_1, x_2, x_3, x_4 be natural numbers, such that $x_1 < x_2 < x_3 < x_4$ and $1+x_1+x_2+x_3+x_4=N$ and the 16 numbers

$$(*) \quad S(1)=1, S(2)=1+x_1, S(3)=1+x_2, S(4)=1+x_1+x_2, \\ S(5)=1+x_3, S(6)=1+x_1+x_3, S(7)=1+x_2+x_3, S(8)=1+x_1+x_2+x_3, \\ S(9)=1+x_4, S(10)=1+x_1+x_4, S(11)=1+x_2+x_4, S(12)=1+x_1+x_2+x_4, \\ S(13)=1+x_3+x_4, S(14)=1+x_1+x_3+x_4, S(15)=1+x_2+x_3+x_4, S(16)=N$$

are pairwise different.

Then the Set S of numbers (*) is a symmetric subset of $\{1, \dots, N\}$.

Let $M=(c_{ij})$, $i, j = 1, \dots, 4$ be a classical magic 4×4 square with entries from $\{1, \dots, 16\}$ and correct rows, columns and diagonals (in the classical sense: every $c_{ij}-1$ is uniquely represented as sum of 1, 2, 4, 8; sufficient: the main diagonal is correct).

The map $k \rightarrow S(k)$ generates a general magic square $M^* = (S(c_{ij}))$, $i, j = 1, \dots, 4$ with magic sum $2(N+1)$

Example

$N=22$, $x_1=3$, $x_2=5$, $x_3=6$, $x_4=7$, $S=\{1, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 22\}$

Take Durerer's magic square for M

16 03 02 13	22 06 04 14
05 10 11 08	07 11 13 15
M: 09 06 07 12	M*: 08 10 12 16
04 15 14 01	09 19 17 01

Proof

Consider any entry c_{ij} of M. Then $c_{ij}-1$ can be uniquely written as a binary 0000, 0001, 0010, 0011, 0100, 0101, 0110, 0111, 1000, 1001, 1010, 1011, 1100, 1101, 1110, 1111.

In M^* read the above binaries as sums of x_1, x_2, x_3, x_4 ; f.i. read 0101 as x_1+x_3 .

Then, because of correctness in any row, column, or diagonal the binary sum of entries $c_{ij}-1$ is $1111+1111=30$,

the sum of entries $S(c_{ij})-1$ in the corresponding row column or diagonal of M^* is $x_4+x_3+x_2+x_4+x_3+x_2+x_1=2N-2$, and therefore, every row, column or diagonal in M^* sums to $2(N+1)$.

Theorem 1.3

Let N be a natural number and let $S=\{s_1, s_2, \dots, s_{16}\}$ be a subset of $\{1, \dots, N\}$ with 16 elements $1=s_1 < s_2 < s_3 < \dots < s_{15} < s_{16}=N$.

Suppose that there exists a general 4x4-magic square with different entries from S , which belongs to the connection figure (1122 3344 5566 7788).

Then there exists a unique set of 4 numbers x_1, x_2, x_3, x_4 with $x_1 < x_2 < x_3 < x_4$ and $1+x_1+x_2+x_3+x_4=N$, such that the 16 numbers (*) of Theorem 1.1 are exactly the members of S .

x_1, x_2, x_3, x_4 can be found as follows:

$x_1=s_2-1$, $x_2=s_3-1$, if $1+x_1+x_2 < s_5$ then $x_3=s_5-1$ else $x_3=s_4-1$, and $x_4=s_{15}-x_2-x_3-1$.

Proof

This follows from (ii) of Theorem 1.1, with $x_1=k, x_2=r, x_3=s$, and $x_4=N-k-r-s-1$.

Remark 1

Every general 4x4 magic square with entry 1 and connection figure (1122 3344 5566 7788) can be generated via $(1, 2, 4, 8) \rightarrow (k, r, s, t)$, $N=k+r+s+t+1$, and the 384 mappings described in (ii) of Theorem 1.1, from only the one classical 4x4-magic square

1	16	4	13
6	11	7	10
15	2	14	3
12	5	9	8.

Remark 2

There is an imbedding map i from the set of general 4x4 magic squares of connection figure (1122 3344 5566 7788) into the set of general 4x4 magic squares of connection figure (1221 3443 5665 7887), namely:

	c01 c02 c03 c04		c01 c05 c06 c02
	c05 c06 c07 c08		c11 c13 c14 c12
i:	c09 c10 c11 c12	-->	c09 c15 c16 c10.
	c13 c14 c15 c16		c03 c07 c08 c04

This is a consequence of four equations, valid for every general 4x4 magic square of connection figure (1122 3344 5566 7788):

$c_{01}+c_{03}+c_{09}+c_{11}=2(N+1)$, $c_{02}+c_{04}+c_{10}+c_{12}=2(N+1)$, $c_{05}+c_{07}+c_{13}+c_{15}=2(N+1)$, $c_{06}+c_{08}+c_{14}+c_{16}=2(N+1)$.